

MODULE 2.4

NONCLINICAL OVERVIEW

Latanoprost/Pharmathen

(eye drops solution containing 50 micrograms/ml (0.005%) latanoprost)

**Pharmacist, Doctor of Medicine,
specialised in Clinical Pharmacology and Toxicology,
Ex-assistant Professor of Forensic Toxicology,
School of Medicine, University of Athens.**

and

**B.Sc., M.Sc., Ph.D. in Clinical Medicine
Senior Chemical & Pharmaceutical Expert Regulatory Affairs
Pharmathen S.A.**

Table of Contents

2.4.1. OVERVIEW OF THE NON-CLINICAL STRATEGY	4
2.4.2. PHARMACOLOGY	4
2.4.2.1. PRIMARY PHARMACODYNAMICS	4
2.4.2.1.1. Mechanism of action.....	4
2.4.2.1.2. Effects on intraocular pressure.....	7
2.4.2.1.3. Effects on blood flow.....	10
2.4.2.1.4. Neuroprotective effects	13
2.4.2.1.5. Effect on platelet-activating factor levels	15
2.4.2.1.6. Additive or synergistic effect.....	15
2.4.2.2. SECONDARY PHARMACODYNAMICS	15
2.4.2.3. SAFETY PHARMACOLOGY	16
2.4.2.4. PHARMACODYNAMIC DRUG INTERACTIONS.....	17
2.4.3. PHARMACOKINETICS.....	18
2.4.3.1. IN VITRO STUDIES	18
2.4.3.2. ANIMAL PHARMACOKINETICS	18
2.4.3.2.1. Pharmacokinetics in the rabbit.....	18
2.4.3.2.2. Pharmacokinetics in the cynomolgus monkey.....	19
2.4.3.2.3. Comparative in-vitro Pharmacokinetic studies performed by Pharmathen	20
2.4.3.3. CLINICAL PHARMACOKINETICS	21
2.4.3.3.1. Ocular pharmacokinetics.....	21
2.4.3.3.2. Systemic pharmacokinetics after intravenous and topical administration on the eye.....	21
2.4.3.3.3. Maximum plasma concentrations during chronic treatment.....	21
2.4.3.3.4. Metabolites in urine and feces	21
2.4.3.3.5. Pharmacokinetics in paediatric patients.....	22
2.4.4. TOXICOLOGY	22
2.4.4.1. SINGLE-DOSE TOXICITY	23
2.4.4.2. REPEAT-DOSE TOXICITY	23
2.4.4.2.1. Topical administration on the eye.....	23
2.4.4.2.2. Oral repeated dose administration	25
2.4.4.2.3. Intravenous repeated dose administration.....	25
2.4.4.3. GENOTOXICITY	26
2.4.4.4. CARCINOGENICITY	26
2.4.4.5. REPRODUCTIVE AND DEVELOPMENT TOXICITY	27
2.4.4.5.1. Fertility and early embryonic development	27
2.4.4.5.2. Embryo-fetal development.....	27
2.4.4.5.3. Prenatal and postnatal development, including maternal function	28
2.4.4.6. LOCAL TOLERANCE	28
2.4.4.6.1. Ocular effects	29

2.4.4.6.1.1. <i>Conjunctival hyperaemia</i>	29
2.4.4.6.1.2. <i>Effects on iris colour and morphology</i>	32
2.4.4.6.1.3. <i>Hypertrichosis and hyperpigmentation of the eyelashes</i>	36
2.4.4.6.1.4. <i>Periocular pigmentation</i>	39
2.4.4.6.1.5. <i>Corneal punctate staining</i>	39
2.4.4.6.1.6. <i>Upper eyelid sulcus deepening</i>	39
2.4.4.6.1.7. <i>Cystoid macular oedema and uveitis</i>	41
2.4.4.6.1.8. <i>Herpes simplex keratitis reactivation</i>	42
2.4.4.6.1.9. <i>Effects on ocular surface</i>	43
2.4.4.6.1.10. <i>Cytotoxic effects</i>	46
2.4.4.6.1.11. <i>Various periocular changes</i>	47
2.4.4.6.1.12. <i>Effects on astrocytes from the lamina cribrosa</i>	47
2.4.4.6.2. <i>Ear disorders</i>	47
2.4.4.7. OTHER TOXICITY STUDIES	47
2.4.4.8. PRESERVATIVE CONSIDERATIONS.....	48
2.4.4.9. DISCUSSION ON EXCIPIENTS	50
2.4.4.10. DISCUSSION ON IMPURITIES.....	51
2.4.5. INTEGRATED OVERVIEW AND CONCLUSIONS	52
2.4.6. LIST OF LITERATURE REFERENCES	59

LIST OF ABBREVIATIONS

AE(s): adverse event(s)

BAK: benzalkonium chloride

BIM: bimatoprost

BP: blood pressure

CAI(s): carbonic anhydrase inhibitor(s)

CCT: central corneal thickness

DOR: dorzolamide

IOP: intraocular pressure

LAT: latanoprost

NTG: normal tension glaucoma

OAG: open-angle glaucoma

ONH: optic nerve head

OH: ocular hypertension

OPP: ocular perfusion pressure

PF: Preservative-free

PG: prostaglandin

PGA(s): prostaglandin analogue(s)

POAG: primary open-angle glaucoma

TIM: timolol

TRAV: travoprost

2.4.1. OVERVIEW OF THE NON-CLINICAL STRATEGY

2.4.6. LIST OF LITERATURE REFERENCES

(Articles in other than English language are included as abstracts,
Very recent articles that are not yet available in our country are included as abstracts)

Abe S, Watabe H, Takaseki S et al (2013): The effects of prostaglandin analogues on intracellular Ca²⁺ in ciliary arteries of wild-type and prostanoid receptor-deficient mice; *J Ocul Pharmacol Ther*, 29, 55-60.

Aihara M, Lindsey JD, Weinreb RN (2002): Reduction of intraocular pressure in mouse eyes treated with latanoprost; *Invest Ophthalmol Vis Sci*, 43, 146-150.

Akaishi T, Kurashima H, Odani-Kawabata N et al (2010): Effects of repeated administrations of tafluprost, latanoprost, and travoprost on optic nerve head blood flow in conscious normal rabbits; *J Ocul Pharmacol Ther*, 26, 181-186.

Albert DM, Gangnon RE, Zimbric ML et al (2004): A study of iridectomy histopathologic features of latanoprost- and non-latanoprost-treated patients; *Arch Ophthalmol*, 122, 1680-1685.

Albert DM, Gangnon RE, Grossniklaus HE et al (2008): A study of histopathological features of latanoprost-treated irides with or without darkening compared with non-latanoprost-treated irides; *Arch Ophthalmol*, 126, 626-31.

Alm A (2014): Latanoprost in the treatment of glaucoma; *Clin Ophthalmol*, 8, 1967-1985.

Alm A, Nilsson SF (2009): Uveoscleral outflow--a review; *Exp Eye Res*, 88, 760-768.

Alm A, Villumsen J (1991): PhXA34, a new potent ocular hypotensive drug: A study on dose-response relationship and on aqueous humor dynamics in healthy volunteers; *Arch Ophthalmol*, 109, 1564-1568.

Alm A, Villumsen J, Tornquist P et al (1993): Intraocular pressure-reducing effect of PhXA41 in patients with increased eye pressure: A one-month study; *Ophthalmology*, 100, 1312-1316.

Alm A, Stjernschantz J, Scandinavian Latanoprost Study Group (1995): Effects on intraocular pressure and side effects of 0.005% latanoprost applied once daily, evening or morning. A comparison with timolol. Scandinavian Latanoprost Study Group; *Ophthalmology*, 102, 1743-1752.

Alvarado JA, Iguchi R, Martinez J et al (2010): Similar effects of selective laser trabeculoplasty and prostaglandin analogs on the permeability of cultured Schlemm canal cells; *Am J Ophthalmol*, 150, 254-264.

Anbar TS, El-Ammawi TS, Barakat M et al (2010): Skin pigmentation after NB-UVB and three analogues of prostaglandin F(2alpha) in guinea pigs: a comparative study; *J Eur Acad Dermatol Venereol*, 24, 28-31.

Anthony TL, Lindsey JD, Weinreb RN (2002): Latanoprost's effects on TIMP-1 and TIMP-2 expression in human ciliary muscle cells; *Invest Ophthalmol Vis Sci*, 43, 3705-3711.

Aptel F, Choudhry R, Stalmans I (2016): Preservative-free versus preserved latanoprost eye drops in patients with open-angle glaucoma or ocular hypertension; *Curr Med Res Opin*, 32, 1457-1463.

Arcieri ES, Santana A, Rocha FN et al (2005): Blood-aqueous barrier changes after the use of prostaglandin analogues in patients with pseudophakia and aphakia: a 6-month randomized trial; *Arch Ophthalmol*, 123, 186-192.

Arend O, Harris A, Wolter P et al (2003): Evaluation of retinal haemodynamics and retinal function after application of dorzolamide, timolol and latanoprost in newly diagnosed open-angle glaucoma patients; *Acta Ophthalmol Scand*, 81, 474-479.

Arranz-Marquez E, Teus MA (2007): Effect of age on the development of a latanoprost-induced increase in iris pigmentation; *Ophthalmology*, 114, 1255-1258.

Arranz-Marquez E, Teus MA (2008): Prostanoids for the management of glaucoma; *Expert Opin Drug Saf*, 7, 801-808.

Arranz-Marquez E, Teus-Guezala MA, Canseco-Gonzalez F (2002): Respiratory effects of chronic therapy with latanoprost in patients with chronic obstructive pulmonary disease; *Arch Soc Esp Ophthalmol*, 77, 553-558.

Arranz-Marquez E, Teus MA, Saornil MA et al (2004): Analysis of irises with a latanoprost-induced change in iris color; *Am J Ophthalmol*, 138, 625-630.

Astin M, Stjernschantz J, Selen G (1994): Role of nitric oxide in PGF₂a-induced ocular hyperemia; *Exp Eye Res*, 59, 401-408.

Ayaki M, Iwasawa A, Inoue Y (2010): Toxicity of antiglaucoma drugs with and without benzalkonium chloride to cultured human corneal endothelial cells; *Clin Ophthalmol*, 4, 1217-1222.

Aydin Kurma S, Acikgoz S, Altun A et al (2014): The effects of topical antiglaucoma drugs as monotherapy on the ocular surface: a prospective study; *J Ophthalmol*, 2014, 460483.

Bafa M, Georgopoulos G, Mihas C et al (2011): The effect of prostaglandin analogues on central corneal thickness of patients with chronic open-angle glaucoma: a 2-year study on 129 eyes; *Acta Ophthalmol*, 89, 448-451.

Bahler CK, Howell KG, Hann CR et al (2008): Prostaglandins increase trabecular meshwork outflow facility in cultured human anterior segments; *Am J Ophthalmol*, 145, 114-119.

Basu S, Sjoquist B (1996): Development of a radioimmunoassay for latanoprost and its application in a long-term study in monkeys; *Prostaglandins Leukot Essent Fatty Acids*, 55, 427-432.

Basu S, Stjernschantz J (1997): Delta13-reductase dependent metabolism of prostaglandins in the mammalian brain and eye; *Prostaglandins Leukot Essent Fatty Acids*, 57, 305-310.

Basu S, Sjoquist B, Stjernschantz J et al (1994): Corneal permeability to and ocular metabolism of phenyl substituted prostaglandin esters in vitro; *Prostaglandins Leukot Essen tatty Acids*, 50, 161-168.

Baudouin C, Riancho L, Warnet JM et al (2007): In vitro studies of antiglaucomatous prostaglandin analogues: travoprost with and without benzalkonium chloride and preserved latanoprost; *Invest Ophthalmol Vis Sci*, 48, 4123-4128.

- Baudouin C, Renard JP, Nordmann JP et al (2012): Prevalence and risk factors for ocular surface disease among patients treated over the long term for glaucoma or ocular hypertension; *Eur J Ophthalmol*, 23, 47-54.
- Bean G, Reardon G, Zimmerman TJ (2004): Association between ocular herpes simplex virus and topical ocular hypotensive therapy; *J Glaucoma*, 13, 361-364.
- Bell NP, Ramos JL, Feldman RM (2010): Safety, tolerability, and efficacy of fixed combination therapy with dorzolamide hydrochloride 2% and timolol maleate 0.5% in glaucoma and ocular hypertension; *Clin Ophthalmol*, 4, 1331.
- Bhattacharjee P, Williams BS, Paterson CA (1999): Responses of intraocular pressure and the pupil of feline eyes to prostaglandin EP1 and FP receptor agonists; *Invest Ophthalmol Vis Sci*, 40, 3047-3053.
- Bolivar G, Sanchez-Barahona C, Teus M et al (2015): Effect of topical prostaglandin analogues on corneal hysteresis; *Acta Ophthalmol* 93, e495-e498.
- Boltz A, Schmidl D, Weigert G et al (2011): Effect of latanoprost on choroidal blood flow regulation in healthy subjects; *Invest Ophthalmol Vis Sci*, 52, 4410-4415.
- Bron A, Chiambaretta F, Pouliquen P et al (2003): Efficacy and safety of substituting a twice-daily regimen of timolol with a single daily instillation of nonpreserved beta-blocker in patients with chronic glaucoma or ocular hypertension; *J Fr Ophthalmol*, 26, 668-674.
- Buguet A, Pv P, Romanet JP (1994): 24-hour (nyctohemeral) and sleep-related variations of intraocular pressure in healthy white individuals; *Am J Ophthalmol*, 117, 342-347.
- Camber O, Edman P (1987): Factors influencing the corneal permeability of prostaglandin F₂ α and its isopropyl ester in vitro; *Int J Pharm*, 37, 27-32.
- Camras CB, The United States Latanoprost Study Group (1996a): Comparison of latanoprost and timolol in patients with ocular hypertension and glaucoma: a six-month masked, multicenter trial in the United States. The United States Latanoprost Study Group; *Ophthalmology*, 103, 138-147.
- Camras CB, Alm A, Watson P et al (1996b): Latanoprost, a prostaglandin analog, for glaucoma therapy. Efficacy and safety after 1 year of treatment in 198 patients. Latanoprost Study Groups; *Ophthalmology*, 103, 1916-1924.
- Camras CB, Wax MB, Ritch R et al (1998): Latanoprost treatment for glaucoma: effects of treating for 1 year and of switching from timolol. United States Latanoprost Study Group; *Am J Ophthalmol*, 126, 390-399.
- Camras CB, Neely DG, Weiss EL (2000): Latanoprost-induced iris color darkening: a case report with long-term follow-up; *J Glaucoma*, 9, 95-98.
- Cardascia N, Vetrugno M, Trabucco T et al (2003): Effects of travoprost eye drops on intraocular pressure and pulsatile ocular blood flow: a 180-day, randomized, double-masked comparison with latanoprost eye drops in patients with open-angle glaucoma; *Curr Ther Res*, 64, 389-400.
- Carvalho AB, Laus JL, Costa VP et al (2006): Effects of travoprost 0.004% compared with latanoprost 0.005% on the intraocular pressure of normal dogs; *Vet Ophthalmol*, 9, 121-125.

Chen CS, Wells J, Craig JE (2004): Topical prostaglandin F(2alpha) analog induced poliosis; *Am J Ophthalmol*, 137, 965-966.

Chen W, Dong N, Huang C et al (2014): Corneal alterations induced by topical application of commercial latanoprost, travoprost and bimatoprost in rabbit; *PLoS One*, 9, e89205.

Chiba T, Kashiwagi K, Kogure S et al (2001): Iridial pigmentation induced by latanoprost ophthalmic solution in Japanese glaucoma patients; *J Glaucoma*, 10, 406-410.

Chiba T, Kashiwagi K, Ishijima K et al (2004): A prospective study of iridial pigmentation and eyelash changes due to ophthalmic treatment with latanoprost; *Jpn J Ophthalmol*, 48, 141-147.

Chiba T, Kashiwagi K, Chiba N et al (2006): Effect of non-steroidal anti-inflammatory ophthalmic solution on intraocular pressure reduction by latanoprost in patients with primary open angle glaucoma or ocular hypertension; *Br J Ophthalmol*, 90, 314-317.

Chou SY, Chou CK, Kuang TM et al (2005): Incidence and severity of iris pigmentation on latanoprost-treated glaucoma eyes; *Eye*, 19, 784-787.

Costagliola C, Prete AD, Incorvaia C et al (2001): Ocular surface changes induced by topical application of latanoprost and timolol: a short-term study in glaucomatous patients with and without allergic conjunctivitis; *Graefes Arch Clin Exp Ophthalmol*, 239, 809-814.

Costagliola C, Parmeggiani F, Virgili G et al (2008): Circadian changes of intraocular pressure and ocular perfusion pressure after timolol or latanoprost in Caucasians with normal-tension glaucoma; *Graefes Arch Clin Exp Ophthalmol*, 246, 389-396.

Costagliola C, dell'Omo R, Romano MR et al (2009): Pharmacotherapy of intraocular pressure - part II. Carbonic anhydrase inhibitors, prostaglandin analogues and prostamides; *Expert Opin Pharmacother*, 10, 2859-2870.

Cracknell KP, Grierson I, Hogg P et al (2003): Latanoprost-induced iris darkening: a morphometric study of human peripheral iridectomies; *Exp Eye Res*, 77, 721-730.

Cracknell KP, Grierson I, Hogg P (2007a): Morphometric effects of long-term exposure to latanoprost; *Ophthalmology*, 114, 938-948.

Cracknell KP, Farnell DJ, Grierson I (2007b): Monte Carlo simulation of latanoprost induced iris darkening; *Comput Methods Programs Biomed*, 87, 93-103.

Crowston JG, Lindsey JD, Aihara M et al (2004a): Effect of latanoprost on intraocular pressure in mice lacking the prostaglandin FP receptor; *Invest Ophthalmol Vis Sci*, 45, 3555-3559.

Crowston JG, Aihara M, Lindsey JD et al (2004b): Effect of latanoprost on outflow facility in the mouse; *Invest Ophthalmol Vis Sci*, 45, 2240-2245.

Cucherat M, Stalmans I, Rouland JF (2014): Relative efficacy and safety of preservative-free latanoprost (T2345) for the treatment of open-angle glaucoma and ocular hypertension: an adjusted Indirect comparison meta-analysis of randomized clinical trials; *J Glaucoma*, 23, e69-e75.

Cvenkel B, Ihan A (2002): Ocular surface changes induced by topical antiglaucoma monotherapy; *Ophthalmologica*, 216, 175-179.

- Daull P, Buggage R, Lambert G et al (2012): A comparative study of a preservative-free latanoprost cationic emulsion (Catioprost) and a BAK-preserved latanoprost solution in animal models; *J Ocul Pharmacol Ther*, 28, 515-523.
- De Santis M, Lucchese A, Carducci B et al (2004): Latanoprost exposure in pregnancy; *Am J Ophthalmol*, 138, 305-306.
- Demirel S, Doganay S, Gurses I et al (2013): Toxic-inflammatory effects of prostoglandin analogs on the ocular surface; *Ocul Immunol Inflamm*, 21, 13-18.
- Demitsu T, Manabe M, Harima N et al (2001): Hypertrichosis induced by latanoprost; *J Am Acad Dermatol*, 44, 721-723.
- Diaz F, Villena A, Vidal L et al (2010): Experimental model of ocular hypertension in the rat: study of the optic nerve capillaries and action of hypotensive drugs; *Invest Ophthalmol Vis Sci*, 51, 946-951.
- Digiuni M, Fogagnolo P, Rossetti L (2012): A review of the use of latanoprost for glaucoma since its launch; *Expert Opin Pharmacother*, 13, 723-745.
- Drago F, Marino A, La Manna C (1999): alpha-Methyl-p-tyrosine inhibits latanoprost-induced melanogenesis in vitro; *Exp Eye Res*, 68, 85-90.
- Drago F, Valzelli S, Emmi I et al (2001): Latanoprost exerts neuroprotective activity in vitro and in vivo; *Exp Eye Res*, 72, 479-486.
- Drance SM, Crichton A, Mills RP (1998): Comparison of the effect of latanoprost 0.005% and timolol 0.5% on the calculated ocular perfusion pressure in patients with normal-tension glaucoma; *Am J Ophthalmol*, 125, 585-592.
- Dutkiewicz R, Albert DM, Levin LA (2000): Effects of latanoprost on tyrosinase activity and mitotic index of cultured melanoma lines; *Exp Eye Res*, 70, 563-569.
- Eisenberg DL, Camras CB (1999): A preliminary risk-benefit assessment of latanoprost and unoprostone in open-angle glaucoma and ocular hypertension; *Drug Saf*, 20, 505-514.
- Elgin U, Batman A, Berker N et al (2006): The comparison of eyelash lengthening effect of latanoprost therapy in adults and children; *Eur J Ophthalmol*, 16, 247-250.
- Emre S, Gul M, Ates B et al (2009): Comparison of the protective effects of prostaglandin analogues in the ischemia and reperfusion model of rabbit eyes; *Exp Anim*, 58, 505-513.
- Erkin EF, Celik P, Kayikcioglu O et al (2006): Effects of latanoprost and betaxolol on cardiovascular and respiratory status of newly diagnosed glaucoma patients; *Ophthalmologica*, 220, 332-337.
- Esaki Y, Shimazaki A, Pellinen P (2016): Ocular tolerability of preservative-free tafluprost and latanoprost: in vitro and in vivo comparative study; *Open Ophthalmol J*, 10, 146.
- Feldman RM (2003): Conjunctival hyperemia and the use of topical prostaglandins in glaucoma and ocular hypertension; *J Ocul Pharmacol Ther*, 19, 23-35.
- Fristrom B, Nilsson SE (1993): Interaction of PhXA41, a new prostaglandin analogue, with pilocarpine: A study on patients with elevated intraocular pressure; *Arch Ophthalmol*, 111, 662-665.

Furuichi M, Chiba T, Abe K et al (2001): Cystoid macular edema associated with topical latanoprost in glaucomatous eyes with a normally functioning blood-ocular barrier; *J Glaucoma*, 10, 233-236.

Gabelt BT, Hennes EA, Bendel MA et al (2009): Prostaglandin subtype-selective and non-selective IOP-lowering comparison in monkeys; *J Ocul Pharmacol Ther*, 25, 1-8.

Gandolfi S, Simmons ST, Sturm R et al (2001): Three-month comparison of bimatoprost and latanoprost in patients with glaucoma and ocular hypertension; *Adv Ther*, 18, 110-121.

Garcia-Feijoo J, Martínez-de-la-Casa JM, Castillo A et al (2006): Circadian IOP-lowering efficacy of travoprost 0.004% ophthalmic solution compared to latanoprost 0.005%; *Curr Med Res Opin*, 22, 1689-1697.

Gelatt KN, MacKay EO (2001): Effect of different dose schedules of latanoprost on intraocular pressure and pupil size in the glaucomatous Beagle; *Vet Ophthalmol*, 4, 283-288.

Georgopoulos GT, Diestelhorst M, Fisher R et al (2002): The short-term effect of latanoprost on intraocular pressure and pulsatile ocular blood flow; *Acta Ophthalmol Scand*, 80, 54-58.

Geyer O, Man O, Weintraub M et al (2001): Acute effect of latanoprost on pulsatile ocular blood flow in normal eyes; *Am J Ophthalmol*, 131, 198-202.

Gherghel D, Hosking SL, Cunliffe IA et al (2008): First-line therapy with latanoprost 0.005% results in improved ocular circulation in newly diagnosed primary open-angle glaucoma patients: a prospective, 6-month, open-label study; *Eye*, 22, 363-369.

Giannico AT, Lima L, Russ HH et al (2013): Eyelash growth induced by topical prostaglandin analogues, bimatoprost, tafluprost, travoprost, and latanoprost in rabbits; *J Ocul Pharmacol Ther*, 29, 817-820.

Gordon YJ, Yates KA, Mah FS et al (2003): The effects of topical Xalatan on the recovery of ocular herpes simplex virus type 1 (HSV-1) in the induced reactivation and spontaneous shedding rabbit models; *J Ocul Pharmacol Ther*, 19, 233-245.

Goto Y, Ibaraki N, Miyake K (2003): Human lens epithelial cell damage and stimulation of their secretion of chemical mediators by benzalkonium chloride rather than latanoprost and timolol; *Arch Ophthalmol*, 121, 835-839.

Grierson I, Lee WR, Albert DM (1999): The fine structure of an iridectomy specimen from a patient with latanoprost-induced eye color change; *Arch Ophthalmol*, 117, 394-396.

Grierson I, Pfeiffer N, Cracknell KP et al (2002): Histology and fine structure of the iris and outflow system following latanoprost therapy; *Surv Ophthalmol*, 47(Suppl.1), S176-S184.

Grierson I, Jonsson M, Cracknell K (2004): Latanoprost and pigmentation; *Jpn J Ophthalmol*, 48, 602-612.

Guenoun JM, Baudouin C, Rat P et al (2005a): In vitro study of inflammatory potential and toxicity profile of latanoprost, travoprost, and bimatoprost in conjunctiva-derived epithelial cells; *Invest Ophthalmol Vis Sci*, 46, 2444-2450.

Guenoun JM, Baudouin C, Rat P et al (2005b): In vitro comparison of cytoprotective and antioxidative effects of latanoprost, travoprost, and bimatoprost on conjunctiva-derived epithelial cells; *Invest Ophthalmol Vis Sci*, 46, 4594-4599.

Gupta SK, Agarwal R, Galpalli ND et al (2007): Comparative efficacy of pilocarpine, timolol and latanoprost in experimental models of glaucoma; *Methods Find Exp Clin Pharmacol*, 29, 665-671.

Hamard P, Blondin C, Debbasch C et al (2003): In vitro effects of preserved and unpreserved antiglaucoma drugs on apoptotic marker expression by human trabecular cells; *Graefes Arch Clin Exp Ophthalmol*, 241, 1037-1043.

Hara (2001): Increased iris pigmentation after use of latanoprost in Japanese brown eyes; *Nippon Ganka Gakkai Zasshi*, 105, 314-321.

Hariharan S, Minocha M, Mishra GP et al (2009): Interaction of ocular hypotensive agents (PGF2 alpha analogs-bimatoprost, latanoprost, and travoprost) with MDR efflux pumps on the rabbit cornea; *J Ocul Pharmacol Ther*, 25, 487-498.

Harris A, Garzosi HJ, McCranor L et al (2009): The effect of latanoprost on ocular blood flow; *Int Ophthalmol*, 29, 19-26.

Hedner J, Svedmyr N, Lunde H et al (1997): The lack of respiratory effects of the ocular hypotensive drug latanoprost in patients with moderate-steroid treated asthma; *Surv Ophthalmol*, 41(Suppl.2), S111-S115.

Hedner J, Everts B, Moller CS (1999): Latanoprost and respiratory function in asthmatic patients: randomized, double-masked, placebo-controlled crossover evaluation; *Arch Ophthalmol*, 117, 1305-1309.

Hernandez M, Urcola JH, Vecino E (2008): Retinal ganglion cell neuroprotection in a rat model of glaucoma following brimonidine, latanoprost or combined treatments; *Exp Eye Res*, 86, 798-806.

Higginbotham EJ, Diestelhorst M, Pfeiffer N et al (2002): The efficacy and safety of unfixed and fixed combinations of latanoprost and other antiglaucoma medications; *Surv Ophthalmol*, 47(Suppl.1), S133-S140.

Hinz B, Rosch S, Ramer R et al (2005): Latanoprost induces matrix metalloproteinase-1 expression in human nonpigmented ciliary epithelial cells through a cyclooxygenase-2-dependent mechanism; *FASEB J*, 19, 1929-1931.

Honda N, Miyai T, Nejima R et al (2010): Effect of latanoprost on the expression of matrix metalloproteinases and tissue inhibitor of metalloproteinase 1 on the ocular surface; *Arch Ophthalmol*, 128, 466-471.

Honrubia F, Garcia-Sanchez J, Polo V et al (2009): Conjunctival hyperaemia with the use of latanoprost versus other prostaglandin analogues in patients with ocular hypertension or glaucoma: a meta-analysis of randomized clinical trials; *Br J Ophthalmol*, 93, 316-321.

Hoyng PF, Rulo AH, Greve EL et al (1997): Fluorescein angiographic evaluation of the effect of latanoprost treatment on blood-retinal barrier integrity: a review of studies conducted on pseudophakic glaucoma patients and on phakic and aphakic monkeys; *Surv Ophthalmol*, 41(Suppl.2), S83-S88.

- Hu DN, Stjernschantz J, McCormick SA (2000): Effect of prostaglandins A(2), E(1), F(2 alpha) and latanoprost on cultured human iridal melanocytes; *Exp Eye Res*, 70, 113-120.
- Husain S, Whitlock NA, Rice DS et al (2006): Effects of latanoprost on rodent intraocular pressure; *Exp Eye Res*, 83, 1453-1458.
- Hylton C, Robin AL (2003): Update on prostaglandin analogs; *Curr Opin Ophthalmol*, 14, 65-69.
- Imesch PD, Wallow IHL, Albert DM (1997): The color of the human eye. A review of morphologic correlates and of some conditions that affect iridial pigmentation; *Surv Ophthalmol*, 41(Suppl.2), S117-S123.
- Inan UU, Ermis SS, Yucel A et al (2003): The effects of latanoprost and brimonidine on blood flow velocity of the retrobulbar vessels: a 3-month clinical trial; *Acta Ophthalmol Scand*, 81, 155-160.
- Inoue K, Wakakura M, Inoue J et al (2006): Adverse reaction after use of latanoprost in Japanese glaucoma patients; *Nippon Ganka Gakkai Zasshi*, 110, 581-587.
- Inoue K, Shiokawa M, Higa R et al (2012): Adverse periocular reactions to five types of prostaglandin analogs; *Eye*, 26, 1465-1472.
- Inoue K, Shiokawa M, Wakakura M et al (2013): Deepening of the upper eyelid sulcus caused by 5 types of prostaglandin analogs; *J Glaucoma*, 22, 626-631.
- Ishii K, Araie M (2000): Effect of topical latanoprost-timolol combined therapy on retinal blood flow and circulation of optic nerve head tissue cynomolgus monkeys; *Jpn J Ophthalmol*, 44, 227-234.
- Ishii K, Tomidokoro A, Nagahara M et al (2001): Effects of topical latanoprost on optic nerve head circulation in rabbits, monkeys, and humans; *Invest Ophthalmol Vis Sci*, 42, 2957-2963.
- Jaenen N, Baudouin C, Pouliquen P et al (2007): Ocular symptoms and signs with preserved and preservative-free glaucoma medications; *Eur J Ophthalmol*, 17, 341-349.
- Jang CH, Cho YB, Choi CH et al (2013): The effect of topically administered latanoprost on the cochlear blood flow and hearing; *Int J Pediatr Otorhinolaryngol*, 77, 981-985.
- Johnson TV, Fan S, Zhan G et al (2010): Efficacy and mechanisms of intraocular pressure reduction with latanoprost and timolol in participants with ocular hypertension: a comparison of 1 and 6 weeks of treatment; *J Glaucoma*, 19, 356-364.
- Johnstone MA (1997): Hypertrichosis and increased pigmentation of eyelashes and adjacent hair in the region of the ipsilateral eyelids of patients treated with unilateral topical latanoprost; *Am J Ophthalmol*, 124, 544-547.
- Johnstone MA, Albert DM (2002): Prostaglandin-induced hair growth; *Surv Ophthalmol*, 47(Suppl.1), S185-S202.
- Jones RL, Woodward DF (2011): Interaction of prostanoid EP₃ and TP receptors in guinea-pig isolated aorta: contractile self-synergism of 11-deoxy-16,16-dimethyl PGE₂; *Br J Pharmacol*, 162, 521-531.
- Kahook MY, Ammar DA (2010): In vitro toxicity of topical ocular prostaglandin analogs and preservatives on corneal epithelial cells; *J Ocul Pharmacol Ther*, 26, 259-263.

Kalouche G, Beguier F, Bakria M et al (2016): Activation of prostaglandin FP and EP2 receptors differently modulates myofibroblast transition in a model of adult primary human trabecular meshwork cells FP and EP2 receptor effects on trabecular meshwork cells; *IOVS*, 57, 1816-1825.

Kanamori A, Naka M, Fukuda M et al (2009): Latanoprost protects rat retinal ganglion cells from apoptosis in vitro and in vivo; *Exp Eye Res*, 88, 535-541.

Karslioglu MZ, Hosal MB, Tekeli O (2015): Periocular changes in topical bimatoprost and latanoprost use; *Turk J Med Sci*, 45, 925-930.

Kashiwagi K, Tsukahara S (2003): Effect of non-steroidal anti-inflammatory ophthalmic solution on intraocular pressure reduction by latanoprost; *Br J Ophthalmol*, 87, 297-301.

Kashiwagi K, Tsukamoto K, Suzuki M et al (2002a): Effects of isopropyl unoprostone and latanoprost on melanogenesis in mouse epidermal melanocytes; *J Glaucoma*, 11, 57-64.

Kashiwagi K, Kanai N, Tsuchida T et al (2002b): Comparison between isopropyl unoprostone and latanoprost by prostaglandin E2 induction, affinity to a prostaglandin transporter, and intra-ocular metabolism; *Exp Eye Res*, 74, 41-49.

Katagiri Y, Takumida M, Hirakawa K et al (2014): Long-term administration of vasopressin can cause Ménière's disease in mice; *Acta Otolaryngol*, 134, 990-1004.

Katz G, Springs CL, Craven ER et al (2010): Ocular surface disease in patients with glaucoma or ocular hypertension treated with either BAK-preserved latanoprost or BAK-free travoprost; *Clin Ophthalmol*, 4, 1253-1261.

Kaufman HE, Varnell ED, Thompson HW (1999): Latanoprost increases the severity and recurrence of herpetic keratitis in the rabbit; *Am J Ophthalmol*, 127, 531-536.

Kaufman HE, Varnell ED, Toshida H et al (2001): Effects of topical unoprostone and latanoprost on acute and recurrent herpetic keratitis in the rabbit; *Am J Ophthalmol*, 131, 643-646.

Kawaguchi I, Higashide T, Ohkubo S et al (2012): Comparison of efficacy of four prostaglandin analogues by bilateral treatment in healthy subjects; *Jpn J Ophthalmol*, 56, 346-353.

Kerstetter JR, Brubaker RF, Wilson SE et al (1988): Prostaglandin F2 α -1-isopropylester lowers intraocular pressure without decreasing aqueous humor flow; *Am J Ophthalmol*, 105, 30-34.

Khoh-Reiter S, Jessen BA (2009): Evaluation of the cytotoxic effects of ophthalmic solutions containing benzalkonium chloride on corneal epithelium using an organotypic 3-D model; *BMC Ophthalmol*, 9, 5.

Kim HJ, Cho BJ (2011): Long-term effect of latanoprost on central corneal thickness in normal tension glaucoma; *J Ocul Pharmacol Ther*, 27, 73-76.

Kim JW, Lindsey JD, Wang N et al (2001): Increased human scleral permeability with prostaglandin exposure; *Invest Ophthalmol Vis Sci*, 42, 1514-1521.

Kim EJ, Kim YH, Kang SH et al (2013): In vitro effects of preservative-free and preserved prostaglandin analogs on primary cultured human conjunctival fibroblast cells; *Korean J Ophthalmol*, 27, 446-453.

Kim JH, Kim EJ, Kim YH et al (2015): In vivo effects of preservative-free and preserved prostaglandin analogs: mouse ocular surface study; *Korean J Ophthalmol*, 29, 270-279.

Kim YK, Na KI, Jeung JW et al (2017): Intraocular pressure-lowering effect of latanoprost is hampered by defective cervical lymphatic drainage; *PLoS One*, 12, e0169683.

Kobayashi H, Kobayashi K (2011): A correlation between latanoprost-induced conjunctival hyperemia and intraocular pressure-lowering effect; *J Glaucoma*, 20, 3-6.

Koh SW, Coll TJ, Rose L et al (2004): Antiglaucoma eye drop pulses--increased interleukin-6 secretion by Tenon's capsule fibroblast cultures; *J Glaucoma*, 13, 200-209.

Konstas AG, Maltezos AC, Gandi S et al (1999): Comparison of 24-hour intraocular pressure reduction with two dosing regimens of latanoprost and timolol maleate in patients with primary open-angle glaucoma; *Am J Ophthalmol*, 128, 15-20.

Koz OG, Ozsoy A, Yarangumeli A et al (2007): Comparison of the effects of travoprost, latanoprost and bimatoprost on ocular circulation: a 6-month clinical trial; *Acta Ophthalmol Scand*, 85, 838-843.

Kozobolis VP, Detorakis ET, Maskaleris G et al (2005): Corneal sensitivity changes following the instillation of latanoprost, bimatoprost, and travoprost eyedrops; *Am J Ophthalmol*, 139, 742-743.

Kraft ME, Glaeser H, Mandery K et al (2010): The prostaglandin transporter OATP2A1 is expressed in human ocular tissues and transports the antiglaucoma prostanoid latanoprost; *Invest Ophthalmol Vis Sci*, 51, 2504-2511.

Kremmer S, Iliadou M, Anastassiou G et al (2014): Influence of latanoprost on retinal microcirculation in glaucoma; *Open Ophthalmol J*, 8, 60-66.

Kuba GB, Kurnatowski-Billion M, Neuser AM et al (2001): Effect of latanoprost on ocular hemodynamics and contrast sensitivity; *Ophthalmologe*, 98, 535-540.

Kucukevcilioglu M, Bayer A, Uysal Y et al (2014): Prostaglandin associated periorbitopathy in patients using bimatoprost, latanoprost and travoprost; *Clin Experiment Ophthalmol*, 42, 126-131.

Kudo H, Nakazawa T, Shimura M et al (2006): Neuroprotective effect of latanoprost on rat retinal ganglion cells; *Graefes Arch Clin Exp Ophthalmol*, 244, 1003-1009.

Kurashima H, Watabe H, Sato N et al (2010): Effects of prostaglandin F(2 α) analogues on endothelin-1-induced impairment of rabbit ocular blood flow: comparison among tafluprost, travoprost, and latanoprost; *Exp Eye Res*, 91, 853-859.

Larsson LI (2001a): Intraocular pressure over 24 hours after single-dose administration of latanoprost 0.005% in healthy volunteers. A randomized, double-masked, placebo controlled, cross-over single center study; *Acta Ophthalmol Scand*, 79, 567-571.

Larsson LI (2001b): Intraocular pressure over 24 hours after repeated administration of latanoprost 0.005% or timolol gel-forming solution 0.5% in patients with ocular hypertension; *Ophthalmology*, 108, 1439-1444.

Larsson LI, Mishima HK, Takamatsu M et al (2002): The effect of latanoprost on circadian intraocular pressure; *Surv Ophthalmol*, 47(Suppl.1), S90-S96.

Lass JH, Eriksson GL, Osterling L et al (2001): Comparison of the corneal effects of latanoprost, fixed combination latanoprost-timolol, and timolol: A double-masked, randomized, one-year study; *Ophthalmology*, 108, 264-271.

Latanoprost-Induced Iris Pigmentation Study Group (2006): Incidence of latanoprost-induced increase in iris pigmentation in Japanese eyes; *Jpn J Ophthalmol*, 50, 96-99.

Lee H, Cho BJ (2015): Long-term effect of latanoprost on central corneal thickness in normal-tension glaucoma: five-year follow-up results; *J Ocul Pharmacol Ther*, 31, 152-155.

Lee S, Kim MK, Choi HJ et al (2013): Comparative cross-sectional analysis of the effects of topical antiglaucoma drugs on the ocular surface; *Adv Ther*, 30, 420-429.

Leng F, Liu P, Li H et al (2011): Long-term topical antiglaucoma medications cause enhanced Tenon's capsule fibroblast proliferation and abnormal TGF- β and MMP expressions: potential effects on glaucoma filtering surgery; *Curr Eye Res*, 36, 301-309.

Li X, He F, B'Ann TG et al (2016): Effects of latanoprost and bimatoprost on the expression of molecules relevant to ocular inflow and outflow pathways; *PloS One*, 11, e0151644.

Liang H, Brignole-Baudouin F, Pauly A et al (2011): Polyquad-preserved travoprost/timolol, benzalkonium chloride (BAK)-preserved travoprost/timolol, and latanoprost/timolol in fixed combinations: a rabbit ocular surface study; *Adv Ther*, 28, 311-325.

Lim KS, Nau CB, O'Byrne MM et al (2008): Mechanism of action of bimatoprost, latanoprost, and travoprost in healthy subjects. A crossover study; *Ophthalmology*, 115, 790.e4-795.e4.

Linden C, Alm A (1997): Effects on intraocular pressure and aqueous flow of various dose regimens of latanoprost in human eyes; *Acta Ophthalmol Scand*, 75, 412-415.

Linden C, Alm A (1998): Latanoprost twice daily is less effective than once daily: indication of receptor subsensitivity?; *Curr Eye Res*, 17, 567-572.

Linden C, Alm A (2001): The effect on intraocular pressure of latanoprost once or four times daily; *Br J Ophthalmol*, 85, 1163-1166.

Linden C, Nuija E, Alm A (1997): Effects on IOP restoration and blood-aqueous barrier after long-term treatment with latanoprost in open angle glaucoma and ocular hypertension; *Br J Ophthalmol*, 81, 370-372.

Lindquist NG, Larsson BS, Stjernschantz J et al (1998): Age-related melanogenesis in the eye of mice, studied by microautoradiography of 3H-methimazole, a specific marker of melanin synthesis; *Exp Eye Res*, 67, 259-264.

Lindquist NG, Larsson BS, Stjernschantz J (1999): Increased pigmentation of iridial melanocytes in primates induced by a prostaglandin analogue; *Exp Eye Res*, 69, 431-436.

Lindsey JD, To HD, Weinreb RN (1994): Induction of c-fos by prostaglandin F2 alpha in human ciliary smooth muscle cells; *Invest Ophthalmol Vis Sci*, 35, 242-250.

Lindsey JD, Jones HL, Hewitt EG et al (2001): Induction of tyrosinase gene transcription in human iris organ cultures exposed to latanoprost; *Arch Ophthalmol*, 119, 853-60.

Liu CJ, Ko YC, Cheng CY et al (2002a): Effect of latanoprost 0.005% and brimonidine tartrate 0.2% on pulsatile ocular blood flow in normal tension glaucoma; *Br J Ophthalmol*, 86, 1236-1239.

Liu CJ, Ko YC, Cheng CY et al (2002b): Changes in intraocular pressure and ocular perfusion pressure after latanoprost 0.005% or brimonidine tartrate 0.2% in normal-tension glaucoma patients; *Ophthalmology*, 109, 2241-2247.

Liu Y, Yanai R, Lu Y et al (2006): Effects of antiglaucoma drugs on collagen gel contraction mediated by human corneal fibroblasts; *J Glaucoma*, 15, 255-259.

Liu Y, Ko JA, Yanai R et al (2008): Induction by latanoprost of collagen gel contraction mediated by human tenon fibroblasts: role of intracellular signaling molecules; *IOVS*, 49, 1429-1436.

Liu JH, Medeiros FA, Slight JR et al (2009): Comparing diurnal and nocturnal effects of brinzolamide and timolol on intraocular pressure in patients receiving latanoprost monotherapy; *Ophthalmology*, 116, 449-454.

Liu Y, Liu Y, Xu D et al (2015): Latanoprost-induced cytokine and chemokine release from human tenon's capsule fibroblasts: role of MAPK and NF- κ B Signaling pathways; *J Glaucoma*, 24, 635-641.

Loeffler KU, Sahm M, Spitznas M (2001): Short-time application of latanoprost does not stimulate melanogenesis in bovine ocular melanin-containing cells in vitro; *Ophthalmic Res*, 33, 102-106.

Lutjen-Drecoll E, Tamm E (1988): Morphological study of the anterior segment of cynomolgus monkey eyes following treatment with prostaglandin F2 alpha; *Exp Eye Res*, 47, 761-769.

Malvitte L, Montange T, Vejux A et al (2007): Measurement of inflammatory cytokines by multicytokine assay in tears of patients with glaucoma topically treated with chronic drugs; *Br J Ophthalmol*, 91, 29-32.

Marchini G, Ghilotti G, Bonadimani M et al (2003): Effects of 0.005% latanoprost on ocular anterior structures and ciliary body thickness; *J Glaucoma*, 12, 295-300.

Maruyama Y, Mori K, Ikeda Y et al (2014): Effects of long-term topical prostaglandin therapy on central corneal thickness; *J Ocul Pharmacol Ther*, Epub ahead of print.

McKibbin M, Menage MJ (1999): The effect of once-daily latanoprost on intraocular pressure and pulsatile ocular blood flow in normal tension glaucoma; *Eye*, 13, 31-34.

Mishima HK, Kiuchi Y, Takamatsu M et al (1997): Circadian intraocular pressure management with latanoprost: diurnal and nocturnal intraocular pressure reduction and increased uveoscleral outflow; *Surv Ophthalmol*, 41(Suppl.2), S139-S144.

Miyake K, Ibaraki N (2002): Prostaglandins and cystoid macular edema; *Surv Ophthalmol*, 47(Suppl.1), S203-S218.

Miyake K, Ota I, Maekubo K et al (1999): Latanoprost accelerates disruption of the blood-aqueous barrier and the incidence of angiographic cystoid macular edema in early postoperative pseudophakias; *Arch Ophthalmol*, 117, 34-40.

Mochizuki H, Itakura H, Yokoyama T et al (2010): Twenty-four-hour ocular hypotensive effects of 0.0015% tafluprost and 0.005% latanoprost in healthy subjects; *Jpn J Ophthalmol*, 54, 286-290.

Moroi SE, Gottfredsdottir MS, Schteingart MT et al (1999): Cystoid macular edema associated with latanoprost therapy in a case series of patients with glaucoma and ocular hypertension; *Ophthalmology*, 106, 1024-1029.

Moschos MM, Nitoda E, Chatziralli IP et al (2016): Impact of prostaglandin glaucoma drops on platelet-activating factor action: an in vitro study; *Drug Des Devel Ther*, 10, 3977-3981.

Nagasubramanian S, Sheth GP, Hitchings RA et al (1993): Intraocular pressure-reducing effect of PhAX41 in ocular hypertension: comparison of dose regimens; *Ophthalmology*, 100, 1305-1311.

Nakakura S, Yamamoto M, Terao E et al (2014): Prostaglandin-associated periorbitopathy in latanoprost users; *Clin Ophthalmol*, 9, 51-56.

Nakanishi Y, Nakamura M, Mukuno H et al (2006): Latanoprost rescues retinal neuro-glial cells from apoptosis by inhibiting caspase-3, which is mediated by p44/p42 mitogen-activated protein kinase; *Exp Eye Res*, 83, 1108-1117.

Nicolela MT, Buckley AR, Walman BE et al (1996a): A comparative study of the effects of timolol and latanoprost on blood flow velocity of the retrobulbar vessels; *Am J Ophthalmol*, 122, 784-789.

Nicolela MT, Walman BE, Buckley AR et al (1996b): Ocular hypertension and primary open-angle glaucoma: a comparative study of their retrobulbar blood flow velocity; *J Glaucoma*, 5, 308-310.

Nilsson SF, Samuelsson M, Bill A et al (1989): Increased uveoscleral outflow as a possible mechanism of ocular hypotension caused by prostaglandin F2 alpha-1-isopropylester in the cynomolgus monkey; *Exp Eye Res*, 48, 707-716.

Nilsson SF, Drecoll E, Lutjen-Drecoll E et al (2006): The prostanoid EP2 receptor agonist butaprost increases uveoscleral outflow in the cynomolgus monkey; *Invest Ophthalmol Vis Sci*, 47, 4042-4049.

Noecker RJ, Herrygers LA, Anwaruddin R (2004): Corneal and conjunctival changes caused by commonly used glaucoma medications; *Cornea*, 23, 490-496.

O'Toole L, Cahill M, O'Brien C (2001): Eyelid hypertrichosis associated with latanoprost is reversible; *Eur J Ophthalmol*, 11, 377-39.

Ocklind A (1998): Effect of latanoprost on the extracellular matrix of the ciliary muscle. A study on cultured cells and tissue sections; *Exp Eye Res*, 67, 179-191.

Oh DJ, Martin JL, Williams AJ et al (2006): Effect of latanoprost on the expression of matrix metalloproteinases and their tissue inhibitors in human trabecular meshwork cells; *Invest Ophthalmol Vis Sci*, 47, 3887-3895.

Ooi YH, Oh DJ, Rhee DJ (2009): Effect of bimatoprost, latanoprost, and unoprostone on matrix metalloproteinases and their inhibitors in human ciliary body smooth muscle cells; *Invest Ophthalmol Vis Sci*, 50, 5259-5265.

Orihashi M, Shima Y, Tsuneki H et al (2005): Potent reduction of intraocular pressure by nipradilol plus latanoprost in ocular hypertensive rabbits; *Biol Pharm Bull*, 28, 65-68.

Orme M, Collins S, Dakin H et al (2010): Mixed treatment comparison and meta-regression of the efficacy and safety of prostaglandin analogues and comparators for primary open-angle glaucoma and ocular hypertension; *Curr Med Res Opin*, 26, 511-528.

Orzalesi N, Rossetti L, Invernizzi T et al (2000): Effect of timolol, latanoprost, and dorzolamide on circadian IOP in glaucoma or ocular hypertension; *Invest Ophthalmol Vis Sci*, 41, 2566-2573.

Orzalesi N, Rossetti L, Bottoli A et al (2003): The effect of latanoprost, brimonidine, and a fixed combination of timolol and dorzolamide on circadian intraocular pressure in patients with glaucoma or ocular hypertension; *Arch Ophthalmol*, 121, 453-457.

Orzalesi N, Rossetti L, Bottoli A et al (2006): Comparison of the effects of latanoprost, travoprost, and bimatoprost on circadian intraocular pressure in patients with glaucoma or ocular hypertension; *Ophthalmology*, 113, 239-246.

Ota T, Aihara M, Narumiya S et al (2005): The effects of prostaglandin analogues on intraocular pressure in prostanoid FP-receptor-deficient mice; *Invest Ophthalmol Vis Sci*, 46, 4159-4163.

Pang IH, Wang WH, Clark AF (2005): Acute effects of glaucoma medications on rat intraocular pressure; *Exp Eye Res*, 80, 207-214.

Park J, Cho HK, Moon JI (2011): Changes to upper eyelid orbital fat from use of topical bimatoprost, travoprost, and latanoprost; *Jpn J Ophthalmol*, 55, 22-27.

Park SH, Yoo SH, Ha SJ (2015): Comparison of ocular pulse amplitude-lowering effects of tafluprost and latanoprost by dynamic contour tonometry; *J Ocul Pharmacol Ther*, 31, 617-622.

Parrish RK, Palmberg P, Sheu WP et al (2003): A comparison of latanoprost, bimatoprost, and travoprost in patients with elevated intraocular pressure: a 12-week, randomized, masked-evaluator multicenter study; *Am J Ophthalmol*, 135, 688-703.

Patel SS, Spencer CM (1996): Latanoprost. A review of its pharmacological properties, clinical efficacy and tolerability in the management of primary open-angle glaucoma and ocular hypertension; *Drugs Aging*, 9, 363-378.

Pauly A, Brignole-Baudouin F, Guenoun JM et al (2007): Comparative study of topical anti-allergic eye drops on human conjunctiva-derived cells: responses to histamine and IFN gamma and toxicological profiles; *Graefes Arch Clin Exp Ophthalmol*, 245, 534-546.

Pauly A, Roubex C, Liang H et al (2012): In vitro and in vivo comparative toxicological study of a new preservative-free latanoprost formulation; *Invest Ophthalmol Vis Sci*, 53, 8172-8180.

Pellinen P, Huhtala A, Tolonen A et al (2012): The cytotoxic effects of preserved and preservative-free prostaglandin analogs on human corneal and conjunctival epithelium in vitro and the distribution of benzalkonium chloride homologs in ocular surface tissues in vivo; *Curr Eye Res*, 37, 145-154.

Perez-Roca F, Rodrigo-Morales E, Garzon I et al (2015): Effects of four formulations of prostaglandin analogs on eye surface cells. A comparative study; *PLoS One*, 10, e0129419.

Perry CM, McGavin JK, Culy CR et al (2003): Latanoprost: an update of its use in glaucoma and ocular hypertension; *Drugs Aging*, 20, 597-630.

- Pfeiffer N, Grierson I, Goldsmith H et al (2001): Histological effects in the iris after 3 months of latanoprost therapy: the Mainz I study; *Arch Ophthalmol*, 119, 191-196.
- Pfeiffer N, Grierson I, Goldsmith H et al (2003): Fine structural evaluation of the iris after unilateral treatment with latanoprost in patients undergoing bilateral trabeculectomy (the Mainz II study); *Arch Ophthalmol*, 121, 23-31.
- Pinheiro R, Panfil C, Schrage N et al (2016): the impact of glaucoma medications on corneal wound healing; *J Glaucoma*, 25, 122-127.
- Poyer JF, Millar C, Kaufman PL (1995): Prostaglandin F2 alpha effects on isolated rhesus monkey ciliary muscle; *Invest Ophthalmol Vis Sci*, 36, 2461-2465.
- Pozarowska D, Pozarowski P, Darzynkiewicz Z (2010): Cytometric assessment of cytostatic and cytotoxic effects of topical glaucoma medications on human epithelial corneal line cells; *Cytometry B Clin Cytom*, 78, 130-137.
- Prota G, Vincensi MR, Napolitano A et al (2000): Latanoprost stimulates eumelanogenesis in iridial melanocytes of cynomolgus monkeys; *Pigment Cell Res*, 13, 147-150.
- Quaranta L, Gandolfo F, Turano R et al (2006): Effects of topical hypotensive drugs on circadian IOP, blood pressure, and calculated diastolic ocular perfusion pressure in patients with glaucoma; *Invest Ophthalmol Vis Sci*, 47, 2917-2923.
- Quaranta L, Miglior S, Floriani I et al (2008): Effects of the timolol-dorzolamide fixed combination and latanoprost on circadian diastolic ocular perfusion pressure in glaucoma; *Invest Ophthalmol Vis Sci*, 49, 4226-4231.
- Raber S, Courtney R, Maeda-Chubachi T et al (2011): Latanoprost systemic exposure in pediatric and adult patients with glaucoma: a phase I, open-label study; *Ophthalmology*, 118, 2022-2027.
- Raber S, Mandema JW, Li H et al (2015): A model-based dose-response meta-analysis of ocular hypotensive agents as a drug development tool to evaluate new therapies in glaucoma; *J Ocul Pharmacol Ther*, 31, 189-197.
- Rabinowitz MP, Katz LJ, Moster MR et al (2015): Unilateral prostaglandin-associated periorbitopathy: a syndrome involving upper eyelid retraction distinguishable from the aging sunken eyelid; *Ophthal Plast Reconstr Surg*, 31, 373-378.
- Racz P, Ruzsonyi MR, Nagy ZT et al (1993): Maintained intraocular pressure reduction with once-a-day application of a new prostaglandin F2 α analogue (PhXA41): An in-hospital, placebo-controlled study; *Arch Ophthalmol*, 111, 657-661.
- Racz P, Ruzsonyi MR, Nagy ZT et al (1996): Around-the-clock intraocular pressure reduction with once-daily application of latanoprost by itself or in combination with timolol; *Arch Ophthalmol*, 114, 268-273.
- Reynolds A, Murray PI, Colloby PS (1998): Darkening of eyelashes in a patient treated with latanoprost; *Eye*, 12, 741-743.
- Richter M, Krauss AH, Woodward DF et al (2003): Morphological changes in the anterior eye segment after long-term treatment with different receptor selective prostaglandin agonists and a prostamide; *Invest Ophthalmol Vis Sci*, 44, 4419-4426.

Ritch R, Ishikawa H, Rothman R et al (2002): The efficacy of latanoprost is independent of the width of the ciliary body face; *J Glaucoma*, 11, 239-243.

Rolle T, Cipullo D, Vizzeri GM et al (2000): Evaluation and comparison between the effects on intraocular pressure and retinal blood flow of two antiglaucomatous drugs administered in monotherapy: brimonidine and latanoprost. Preliminary results; *Acta Ophthalmol Scand Suppl*, 232, 50-52.

Rouland JF, Traverso CE, Stalmans I et al (2013): Efficacy and safety of preservative-free latanoprost eyedrops, compared with BAK-preserved latanoprost in patients with ocular hypertension or glaucoma; *Br J Ophthalmol*, 97, 196-200.

Rulo AH, Greve EL, Hoyng PF (1994): Additive effect of latanoprost, a prostaglandin F2 alpha analogue, and timolol in patients with elevated intraocular pressure; *Br J Ophthalmol*, 78, 899-902.

Russo A, Riva I, Pizzolante T et al (2008): Latanoprost ophthalmic solution in the treatment of open angle glaucoma or raised intraocular pressure: a review; *Clin Ophthalmol*, 2, 897-905.

Sagara T, Gatton DD, Lindsey JD et al (1999): Topical prostaglandin F2alpha treatment reduces collagen types I, III, and IV in the monkey uveoscleral outflow pathway; *Arch Ophthalmol*, 117, 794-801.

Sakata R, Shirato S, Miyata K et al (2013): Recovery from deepening of the upper eyelid sulcus after switching from bimatoprost to latanoprost; *Jpn J Ophthalmol*, 57, 179-184.

Sanford M (2014): Preservative-free latanoprost eye drops in patients with primary open-angle glaucoma/ocular hypertension; *Clin Drug Investig*, 34, 521-528.

Schumer RA, Camras CB, Mandahl AK (2002): Putative side effects of prostaglandin analogs; *Surv Ophthalmol*, 47(Suppl.1), S219-S230.

Selen G, Stjernschantz J, Resul B (1997): Prostaglandin-induced iridial pigmentation in primates; *Surv Ophthalmol*, 41(Suppl.2), S125-S128.

Selen F, Tekeli O, Yanik O (2017): Assessment of the anterior chamber flare and macular thickness in patients treated with topical antiglaucomatous drugs; *J Ocul Pharmacol Ther*, Epub ahead of print.

Sen E, Nalcacioglu P, Yazici A et al (2008): Comparison of the effects of latanoprost and bimatoprost on central corneal thickness; *J Glaucoma*, 17, 398-402.

Seong GJ, Lee HK, Hong YJ (1999): Effects of 0.005% latanoprost on optic nerve head and peripapillary retinal blood flow; *Ophthalmologica*, 213, 355-359.

Serle JB, Wang RF, Mittag TW et al (2001): Effect of pilocarpine 4% in combination with latanoprost 0.005% or 8-iso prostaglandin E2 0.1% on intraocular pressure in laser-induced glaucomatous monkey eyes; *J Glaucoma*, 10, 215-219.

Sharif NA, Crider JY, Husain S et al (2003a): Human ciliary muscle cell responses to FP-class prostaglandin analogs: phosphoinositide hydrolysis, intracellular Ca²⁺ mobilization and MAP kinase activation; *J Ocul Pharmacol Ther*, 19, 437-455.

Sharif NA, Kelly CR, Crider JY et al (2003b): Ocular hypotensive FP prostaglandin (PG) analogs: PG receptor subtype binding affinities and selectivities, and agonist potencies at FP and other PG receptors in cultured cells; *J Ocul Pharmacol Ther*, 19, 501-515.

Sharif NA, Kelly CR, Crider JY (2003c): Human trabecular meshwork cell responses induced by bimatoprost, travoprost, unoprostone, and other FP prostaglandin receptor agonist analogues; *Invest Ophthalmol Vis Sci*, 44, 715-721.

Sharpe ED, Reynolds AC, Skuta GL et al (2007): The clinical impact and incidence of periocular pigmentation associated with either latanoprost or bimatoprost therapy; *Curr Eye Res*, 32, 1037-1043.

Shen JW, Shan M, Peng YY et al (2016): Cytotoxic effect of latanoprost on human corneal stromal cells in vitro and its possible mechanisms; *Curr Eye Res*, *Epub ahead of print*.

Shin JA, Kim NW, Kang SS et al (2016): In vitro effects of prostaglandin analogs on cultured astrocytes obtained from the lamina cribrosa; *Curr Eye Res*, 41, 676-682.

Sjoquist B, Stjernschantz J (2002): Ocular and systemic pharmacokinetics of latanoprost in humans; *Surv Ophthalmol*, 47(Suppl.1), S6-S12.

Sjoquist B, Basu S, Byding P et al (1998): The pharmacokinetics of a new antiglaucoma drug, latanoprost, in the rabbit; *Drug Metab Dispos*, 26, 745-754.

Sjoquist B, Tajallaei S, Stjernschantz J (1999a): Pharmacokinetics of latanoprost in the cynomolgus monkey. 1st communication: single intravenous, oral or topical administration on the eye; *Arzneimittelforschung*, 49, 225-233.

Sjoquist B, Uhlin A, Byding P et al (1999b): Pharmacokinetics of latanoprost in the cynomolgus monkey. 2nd communication: repeated topical administration on the eye; *Arzneimittelforschung*, 49, 234-239.

Sjoquist B, Johansson A, Stjernschantz J (1999c): Pharmacokinetics of latanoprost in the cynomolgus monkey. 3rd communication: tissue distribution after topical administration on the eye studied by whole body autoradiography. Glaucoma Research Laboratories; *Arzneimittelforschung*, 49, 240-249.

Smedowski A, Paterno JJ, Toropainen E et al (2014): Excipients of preservative-free latanoprost induced inflammatory response and cytotoxicity in immortalized human HCE-2 corneal epithelial cells; *J Biochem Pharmacol Res*, 2, 175.

Smith-Thomas L, Moustafa M, Spada CS et al (2004): Latanoprost-induced pigmentation in human iridial melanocytes is fibroblast dependent; *Exp Eye Res*, 78, 973-985.

Sponsel WE, Paris G, Trigo Y et al (2002a): Latanoprost and brimonidine: therapeutic and physiologic assessment before and after oral nonsteroidal anti-inflammatory therapy; *Am J Ophthalmol*, 133, 11-18.

Sponsel W, Paris G, Trigo Y et al (2002b): Comparative effects of latanoprost (Xalatan) and unoprostone (Rescula) in patients with open-angle glaucoma and suspected glaucoma; *Am J Ophthalmol*, 134, 552-557.

Stewart WC, Kolker AE, Stewart JA et al (2003a): Conjunctival hyperemia in healthy subjects after short-term dosing with latanoprost, bimatoprost, and travoprost; *Am J Ophthalmol*, 135, 314-320.

Stewart WC, Stewart JA, Jenkins JN et al (2003b): Corneal punctate staining with latanoprost, bimatoprost, and travoprost in healthy subjects; *J Glaucoma*, 12, 475-479.

Stjernschantz JW (2001): From PGF(2alpha)-isopropyl ester to latanoprost: a review of the development of xalatan: the Proctor Lecture; *Invest Ophthalmol Vis Sci*, 42, 1134-1145.

Stjernschantz JW, Alm A (1996): Latanoprost as a new horizon in the medical management of glaucoma; *Curr Opin Ophthalmol*, 7, 11-17.

Stjernschantz J, Selen G, Sjoquist B et al (1995): Preclinical pharmacology of latanoprost, a phenyl-substituted PGF2 alpha analogue; *Adv Prostaglandin Thromboxane Leukot Res*, 23, 513-518.

Stjernschantz J, Selen G, Astin M et al (1999): Effect of latanoprost on regional blood flow and capillary permeability in the monkey eye; *Arch Ophthalmol*, 117, 1363-1367.

Stjernschantz J, Selen G, Astin M et al (2000): Microvascular effects of selective prostaglandin analogues in the eye with special reference to latanoprost and glaucoma treatment; *Prog Retin Eye Res*, 19, 459-496.

Stjernschantz JW, Albert DM, Hu DN et al (2002): Mechanism and clinical significance of prostaglandin-induced iris pigmentation; *Surv Ophthalmol*, 47(Suppl.1), S162-S175.

Strober BE, Potash S, Grossman ME (2001): Eyelash hypertrichosis in a patient treated with topical latanoprost; *Cutis*, 67, 109-110.

Studer ME, Martin CL, Stiles J (2000): Effects of 0.005% latanoprost solution on intraocular pressure in healthy dogs and cats; *Am J Vet Res*, 61, 1220-1224.

Sudesh S, Cohen EJ, Rapuano CJ et al (1999): Corneal toxicity associated with latanoprost; *Arch Ophthalmol*, 117, 539-540.

Sugiyama T, Kojima S, Ishida O et al (2009): Changes in optic nerve head blood flow induced by the combined therapy of latanoprost and beta blockers; *Acta Ophthalmol*, 87, 797-800.

Taketani Y, Yamagishi R, Fujishiro T et al (2014): Activation of the prostanoid FP receptor inhibits adipogenesis leading to deepening of the upper eyelid sulcus in prostaglandin-associated periorbitopathy; *Invest Ophthalmol Vis Sci*, 55, 1269-1276.

Tamaki Y, Nagahara M, Araie M et al (2001): Topical latanoprost and optic nerve head and retinal circulation in humans; *J Ocul Pharmacol Ther*, 17, 403-411.

Tamer C, Oksuz H (2007): Circadian intraocular pressure control with dorzolamide versus timolol maleate add-on treatments in primary open-angle glaucoma patients using latanoprost; *Ophthalmic Res*, 39, 24-31.

Tamm E, Rittig M, Lutjen-Drecoll E (1990): Electron microscopy and immunohistochemical studies of the intraocular pressure lowering effect of prostaglandin F2 alpha; *Fortschr Ophthalmol*, 87, 623-629.

- Terai N, Schlotzer-Schrehardt U, Lampel J et al (2009): Effect of latanoprost and timolol on the histopathology of the human conjunctiva; *Br J Ophthalmol*, 93, 219-224.
- Teus MA, Arranz-Márquez E, Lucea-Suescun P (2002): Incidence of iris colour change in latanoprost treated eyes; *Br J Ophthalmol*, 86, 1085-1088.
- Thygesen J, Aaen K, Theodorsen F et al (2000): Short-term effect of latanoprost and timolol eye drops on tear fluid and the ocular surface in patients with primary open-angle glaucoma and ocular hypertension; *Acta Ophthalmol Scand*, 78, 37-44.
- Toris CB, Camras CB, Yablonski ME (1993): Effects of PhXA41, a new prostaglandin F₂ α analog, on aqueous humor dynamics in human eyes; *Ophthalmology*, 100, 1297-1304.
- Tressler CS, Wiseman RL, Dombi TM et al (2011): Lack of evidence for a link between latanoprost use and malignant melanoma: an analysis of safety databases and a review of the literature; *Br J Ophthalmol*, 95, 1490-1495.
- Trzeciacka A, Paterno JJ, Toropainen E et al (2016): Long-term topical application of preservative-free prostaglandin analogues evokes macrophage infiltration in the ocular adnexa; *Eur J Pharmacol*, 788, 12-20.
- Tsai JC, Sivak-Callcott JA, Haik BG et al (2001): Latanoprost-induced iris heterochromia and open-angle glaucoma: a clinicopathologic report; *J Glaucoma*, 10, 411-413.
- Tsikripis P, Papaconstantinou D, Koutsandrea C et al (2013): The effect of prostaglandin analogs on the biomechanical properties and central thickness of the cornea of patients with open-angle glaucoma: a 3-year study on 108 eyes; *Drug Des Devel Ther*, 7, 1149-1156.
- Uematsu M, Mohamed YH, Onizuka N et al (2016): Acute corneal toxicity of latanoprost with different preservatives; *Cutan Ocul Toxicol*, 35, 120-125.
- Uno H, Zimbric ML, Albert DM et al (2002): Effect of latanoprost on hair growth in the bald scalp of the stump-tailed macaque: a pilot study; *Acta Derm Venereol*, 82, 7-12.
- Van Alphen GWHM, Wilhelm PB, Elsenfeld PW (1977): The effect of prostaglandins on the isolated internal muscles of the mammalian eye, including man; *Doc Ophthalmol*, 42, 397-415.
- Vetrugno M, Cantatore F, Gigante G et al (1998): Latanoprost 0.005% in POAG: effects on IOP and ocular blood flow; *Acta Ophthalmol Scand Suppl*, 227, 40-41.
- Vidal L, Diaz F, Villena A et al (2010): Reaction of Müller cells in an experimental rat model of increased intraocular pressure following timolol, latanoprost and brimonidine; *Brain Res Bull*, 82, 18-24.
- Villumsen J, Alm A (1989): Prostaglandin F₂ α -isopropylester eye drops: effects in normal human eyes; *Br J Ophthalmol*, 73, 419-426.
- Walimbe T, Chelerkar V, Bhagat P et al (2016): Effect of benzalkonium chloride-free latanoprost ophthalmic solution on ocular surface in patients with glaucoma; *Clin Ophthalmol*, 10, 821-827.
- Wand M (1997): Latanoprost and hyperpigmentation of eyelashes; *Arch Ophthalmol*, 115, 1206-1208.

- Wand M, Gaudio AR, Shields MB (2001): Latanoprost and cystoid macular edema in high-risk aphakic or pseudophakic eyes; *J Cataract Refract Surg*, 27, 1397-1401.
- Wang RF, Podos SM, Serle JB et al (2000): Effect of latanoprost or 8-iso prostaglandin E2 alone and in combination on intraocular pressure in glaucomatous monkey eyes; *Arch Ophthalmol*, 118, 74-77.
- Wang NL, Lu QJ, Li JH et al (2008): Prostaglandin induces the expression of matrix metalloproteinase-1 in ciliary melanocytes; *Chin Med J (Engl)*, 121, 1173-1176.
- Wang G, Xu X, Yao X et al (2013): Latanoprost effectively ameliorates glucose and lipid disorders in db/db and ob/ob mice; *Diabetologia*, 56, 2702-2712.
- Warwar RE, Bullock JD, Ballal D (1998): Cystoid macular edema and anterior uveitis associated with latanoprost use; Experience and incidence in a retrospective review of 94 patients; *Ophthalmology*, 105, 263-268.
- Watson P, Stjernschantz J, The Latanoprost Study Group (1996): A six-month, randomized, double-masked study comparing latanoprost with timolol in open-angle glaucoma and ocular hypertension. The Latanoprost Study Group; *Ophthalmology*, 103, 126-137.
- Watson PG, the Latanoprost Study Group (1998): Latanoprost: Two years' experience of its use in the United Kingdom; *Ophthalmology*, 105, 82-87.
- Weinreb RN, Lindsey JD (2002): Metalloproteinase gene transcription in human ciliary muscle cells with latanoprost; *Invest Ophthalmol Vis Sci*, 43, 716-722.
- Weinreb RN, Kashiwagi K, Kashiwagi F et al (1997): Prostaglandins increase matrix metalloproteinase release from human ciliary smooth muscle cells; *Invest Ophthalmol Vis Sci*, 38, 2772-2780.
- Weinreb RN, Lindsey JD, Marchenko G et al (2004): Prostaglandin FP agonists alter metalloproteinase gene expression in sclera; *Invest Ophthalmol Vis Sci*, 45, 4368-4377.
- Wierzbowska J, Lagocka A, Siemiatkowska A et al (2009): The impact of common classes of topical antiglaucoma medications on central corneal thickness--own observations; *Klin Oczna*, 111, 323-326.
- Wistrand PJ, Stjernschantz J, Olsson K (1997): The incidence and time-course of latanoprost-induced iridial pigmentation as a function of eye color; *Surv Ophthalmol*, 41(Suppl.2), S129-S138.
- Wu KY, Wu PC, Lai YH et al (2013): Novel usage of intraocular pressure-lowering drugs as wound-healing inhibitors after trabeculectomy with cell culture and animal models; *Kaohsiung J Med Sci*, 29, 353-361.
- Wu N, Chen Y, Yu X et al (2016): Changes in corneal biomechanical properties after long-term topical prostaglandin therapy; *PLoS One*, 11, e0155527.
- Xalatan Product Monograph (2014): Latanoprost Ophthalmic Solution, 50 µg/mL; Assesed online at: http://www.pfizer.ca/sites/g/files/g10017036/f/201410/Xalatan_0.pdf [Accessed on 3 February 2017].

Xalatan SPC (2017): Xalatan 0.005% eye drops solution; Accessed online at: <http://www.medicines.org.uk/emc/medicine/9043/SPC/Xalatan+0.005++eye+drops+solution/> [Accessed on 14 February 2017].

Yamada H, Yoneda M, Goshō M et al (2016): Bimatoprost, latanoprost, and tafluprost induce differential expression of matrix metalloproteinases and tissue inhibitor of metalloproteinases; *BMC Ophthalmol*, 16, 26.

Yamagishi R, Aihara M, Araie M (2011): Neuroprotective effects of prostaglandin analogues on retinal ganglion cell death independent of intraocular pressure reduction; *Exp Eye Res*, 93, 265-270.

Yamaji K, Yoshitomi T, Ishikawa H et al (2005): Prostaglandins E1 and E2, but not F2alpha or latanoprost, inhibit monkey ciliary muscle contraction; *Curr Eye Res*, 30, 661-665.

Yamamoto K, Sato K, Yukita M et al (2017): The neuroprotective effect of latanoprost acts via klotho-mediated suppression of calpain activation after optic nerve transection; *J Neurochem*, 140, 495-508.

Yanagi M, Kiuchi Y, Yuasa Y et al (2016): Association between glaucoma eye drops and hyperemia; *Jpn J Ophthalmol*, 60, 72-77.

Yeh PC, Ramanathan S (2002): Latanoprost and clinically significant cystoid macular edema after uneventful phacoemulsification with intraocular lens implantation; *J Cataract Refract Surg*, 28, 1814-1818.

Yoo R, Choi YA, Cho BJ (2017): Change in central corneal thickness after the discontinuation of latanoprost in normal tension glaucoma-change in central corneal thickness after stop of latanoprost; *J Ocul Pharmacol Ther*, 33, 57-61.

You JY, Cho BJ (2013): Effect of latanoprost on central corneal thickness in unilateral normal-tension glaucoma; *J Ocul Pharmacol Ther*, 29, 335-338.

Yousufzai SY, Ye Z, Abdel-Latif AA (1996): Prostaglandin F2 alpha and its analogs induce release of endogenous prostaglandins in iris and ciliary muscles isolated from cat and other mammalian species; *Exp Eye Res*, 63, 305-310.

Zeitz O, Matthiessen ET, Reuss J et al (2005): Effects of glaucoma drugs on ocular hemodynamics in normal tension glaucoma: a randomized trial comparing bimatoprost and latanoprost with dorzolamide; *BMC Ophthalmology*, 5, 1-11.

Zhan GL, Toris CB, Camras CB et al (1998): Prostaglandin-induced iris color darkening; *Arch Ophthalmol*, 116, 1065-1068.

Zhao X, Pearson KE, Stephan DA et al (2003): Effects of prostaglandin analogues on human ciliary muscle and trabecular meshwork cells; *Invest Ophthalmol Vis Sci*, 44, 1945-1952.

Zheng J, Feng X, Hou L et al (2011): Latanoprost promotes neurite outgrowth in differentiated RGC-5 cells via the PI3K-Akt-mTOR signaling pathway; *Cell Mol Neurobiol*, 31, 597-604.

Zhong Y, Shen X, Yu J et al (2011): The comparison of the effects of latanoprost, travoprost, and bimatoprost on central corneal thickness; *Cornea*, 30, 861-864.

Ziai N, Dolan JW, Kacere RD et al (1993): The effects on aqueous dynamics of PhXA41, a new prostaglandin F2 α analogue, after topical application in normal and ocular hypertensive human eyes; *Arch Ophthalmol*, 111, 1351-1358.